

Linz, June 12, 2000

Critical light scattering in pure liquids¹

G. Flossmann² and R. Folk^{2,3}

¹ Paper presented at the Fourteenth Symposium on Thermophysical Properties,
June 25-30, 2000, Boulder, Colorado, U.S.A.

²Institute for Theoretical Physics, University of Linz, Austria

³To whom correspondence should be addressed.

ABSTRACT

We review light scattering experiments near the critical point (T_c, ρ_c) in fluid systems in particular the central Rayleigh peak in the frequency spectrum. Within a nonasymptotic renormalization group theory we calculate the crossover function between several regions: (i) from the background to the asymptotic region, (ii) from the hydrodynamic region (wave length \gg correlation length) to the critical region (wave length \ll correlation length), (iii) from critical densities to noncritical densities. Contrary to the mode coupling expression the appropriate scaling function is well defined in all limits of its arguments. At T_c we also consider the crossover in the wave vector dependence of the linewidth. We then compare our theoretical result with experiments pure liquids. In doing so we choose the nonuniversal parameters in consistency with the transport coefficients (i.e. the shear viscosity) for the same substance which can be evaluated within the same formalism.

KEY WORDS: characteristic frequency; critical point; dynamic critical phenomena; light scattering; renormalization group theory; transport coefficients

1. INTRODUCTION

Light scattering is a common technique to investigate the transport properties of a fluid system. Especially the characteristic frequency ω_c , defined as the half width at half height of the central Rayleigh peak in the frequency spectrum, provides useful information about the dynamical properties of the system.

Far away from the critical point in the background region, characterized by small values of the correlation length ξ or large values of the wave vector k , the behavior of the characteristic frequency is described by the van Hove theory. In the hydrodynamic region for small values of $k\xi$ the characteristic frequency is given by $\omega_c = D_T(T, \rho) k^2$ with $D_T(T, \rho)$ being the regular thermal diffusivity (the ratio of the nonsingular thermal conductivity and the specific heat at constant pressure), whereas in the 'background' critical region, for large values of $k\xi$, the characteristic frequency is proportional to k^4 .

In the vicinity of the critical point, characterized by large values of ξ or small values of k , fluctuations determine the form of the characteristic frequency. In the critical limit for large values of $k\xi$ the power law behavior $\omega_c \sim k^z$ with the dynamical critical exponent $z \approx 3$ is expected. In the hydrodynamic limit for $k\xi \rightarrow 0$ the characteristic frequency is again given by $\omega_c = D_T(T, \rho) k^2$ but now with a singular Onsager coefficient and thus a diverging thermal conductivity.

Within the nonasymptotic renormalization group theory we are able to describe the crossover from the asymptotic region to the background region as well as from the hydrodynamic region to the region of large wave vectors quantitatively and compare our theory to experiments in pure liquids.

2. THE CHARACTERISTIC FREQUENCY

The characteristic frequency ω_c can be found from the dynamic order parameter correlation function within the dynamic model H, which is shortly reviewed in Ref. [1]. In one-loop perturbation theory the characteristic frequency can be found from the order parameter vertex function so that after renormalization ω_c , expressed in terms of the renormalized model parameters $\Gamma(\ell)$ and $f_t(\ell)$, is found to be of the form

$$\omega_c(k, \xi) = \Gamma(\ell) k^2 (\xi^{-2} + k^2) \left\{ 1 - \frac{f_t^2(\ell)}{16} \left[-5 + 6 x^{-2} \ln(1 + x^2) \right] \right\} \quad (1)$$

with $x = k\xi$. The temperature dependence enters via the flow equations for the mode coupling $f_t(\ell)$ and the Onsager coefficient $\Gamma(\ell)$, which are given by Eqs. (7) and (8) of Ref. [1]. Other than for the evaluation of the frequency dependent shear viscosity the connection between the flow parameter ℓ and the correlation length or the wave vector respectively is found from the matching condition (which arises naturally within the calculation of the vertex function)

$$\left(\xi_0^{-1} \ell \right)^2 = \xi^{-2} + k^2, \quad (2)$$

where ξ_0 is the amplitude in the power law for the correlation length $\xi = \xi_0 t^{-\nu}$. As described in Ref. [1] the correlation length may also be expressed in terms of the reduced density using the cubic model.

In order to discuss the various limits of the characteristic frequency it is useful to rewrite Eq. (1): Extracting the critical asymptotics the characteristic frequency reads

$$\omega_c(k, x) = \Gamma_{as} k^z \left(\frac{1 + x^2}{x^2} \right)^{1-x_\lambda/2} [c_{na}(k, x)]^{x_\lambda} f(k, x) \quad (3)$$

with $x = k\xi$, $z = 4 - x_\lambda$. The critical exponent of the thermal conductivity x_λ is related to the exponent of the shear viscosity x_η by $x_\lambda + x_\eta = 1$ in one loop order. The function $f(k, x)$ defined as (f_t^{*2} is the fixed point value of the mode coupling)

$$f(k, x) = 1 - \frac{f_t^{*2}}{16 c_{na}(k, x)} \left[-5 + 6 x^{-2} \ln(1 + x^2) \right]. \quad (4)$$

The nonasymptotic contributions are collected in

$$c_{na}(k, x) = \left[1 + \frac{k}{k_0} \sqrt{\frac{1 + x^2}{x^2}} \right] \quad (5)$$

so that the asymptotic region is characterized by $c_{na}(k, x) = 1$ and there the frequency can indeed be written in scaling form with the scaling function f depending only on the variable $x = k\xi$. And finally the asymptotic Onsager coefficient Γ_{as} and the crossover wave length k_0 are given by

$$\Gamma_{as} = \Gamma_0 \left(\frac{f_0^2 \ell_0}{f_t^{*2} \xi_0} \right)^{x_\lambda}, \quad k_0^{-1} = \left(\frac{f_t^{*2}}{f_0^2} - 1 \right) \frac{\xi_0}{\ell_0} \quad (6)$$

containing the nonuniversal initial values Γ_0 , and f_0 at ℓ_0 corresponding to a temperature t_0 .

3. CROSSOVER BEHAVIOR

We start this section with a discussion of the various limits of the characteristic frequency: In the *hydrodynamic* limit $x = k\xi \rightarrow 0$ we find

$$\lim_{x \rightarrow 0} \omega_c(k, \xi) = \Gamma_{as} k^2 \xi^{-2+x_\lambda} \left[1 + \frac{1}{x_0} \right]^{x_\lambda} \left\{ 1 - \frac{f_t^{*2}}{16} \left[1 + \frac{1}{x_0} \right]^{-1} \right\}, \quad (7)$$

with $x_0 = k_0 \xi$. Here the coefficient of k^2 is the nonasymptotic expression for the temperature dependent thermal diffusion coefficient $D_T(\xi)$ discussed in Ref. [2]

so that we can rewrite Eq. (7) in the well-known form $\omega_c = D_T(\xi)k^2$ for the hydrodynamic region. In the opposite *critical* limit $x \rightarrow \infty$ we have

$$\lim_{x \rightarrow \infty} \omega_c(k, \xi) = \omega_c(k) = \Gamma_{as} k^z \left[1 + \frac{k}{k_0} \right]^{x_\lambda} \left\{ 1 + \frac{5f_t^{*2}}{16} \left[1 + \frac{k}{k_0} \right]^{-1} \right\}, \quad (8)$$

which is the wave vector dependent nonasymptotic expression of the characteristic frequency. Both nonasymptotic expressions allow to discuss the crossover from the *asymptotic* limit $\xi k_0 \rightarrow \infty$ or $k/k_0 \rightarrow 0$ to the *background* limit $\xi k_0 \rightarrow 0$ or $k/k_0 \rightarrow \infty$ respectively. In the hydrodynamic case we obtain the limits

$$\lim_{\xi k_0 \rightarrow \infty} \omega_c(k, \xi) = \Gamma_{as} k^2 \xi^{-2+x_\lambda} \left(1 - \frac{f_t^{*2}}{16} \right), \quad (9)$$

$$\lim_{\xi k_0 \rightarrow 0} \omega_c(k, \xi) = \Gamma_0 k^2 \xi^{-2} \left(1 - \frac{f_0^2}{f_t^{*2}} \right)^{x_\lambda}, \quad (10)$$

where we used the expression for k_0 given in Eq.(6) for the last limit. In the background limit our expression reaches the van Hove behavior $\omega_c \propto k^2 \xi^{-2}$. In the critical region we obtain

$$\lim_{k/k_0 \rightarrow 0} \omega_c(k) = \Gamma_{as} k^z \left(1 + \frac{5f_t^{*2}}{16} \right), \quad (11)$$

$$\lim_{k/k_0 \rightarrow \infty} \omega_c(k) = \Gamma_0 k^4 \left(1 - \frac{f_0^2}{f_t^{*2}} \right)^{x_\lambda}, \quad (12)$$

where again we reach the van Hove theory for large values of the ratio k/k_0 . This means that our results for the characteristic frequency describe the crossover in the correlation length (from ξ^{-2+x_λ} to ξ^{-2}) in the hydrodynamic region characterized by the limit $x \rightarrow 0$ as well as the crossover in the wave vector (from k^z to k^4) in the critical region characterized by the limit $x \rightarrow \infty$. This is shown in Fig. 1 where we have plotted the dimensionless characteristic frequency $\omega_c/\Gamma_0 k^2$ as function of the wave vector and the correlation length along the critical isochore. In

Fig. 2 we have plotted $\omega_c/\Gamma_0 k^2$ as a function of the reduced temperature for various values of the reduced density and the wave vector using the cubic model described in Ref. [1] (the density enters via the correlation length in the matching condition). Only at zero wave vector and along the critical isochore (that means for $\Delta\rho = 0$) we can see the asymptotic power-law behavior $\omega_c \sim \xi^{-2+x_\lambda} \sim t^{\nu(2-x_\lambda)}$ and the crossover to the van Hove theory with $\omega_c \sim t^{2\nu}$ whereas for noncritical densities or non vanishing wave vector the dimensionless characteristic frequency approaches a finite value for $t \rightarrow 0$.

We have seen that with our nonasymptotic theory we always reach the van Hove behavior in the nonasymptotic limit for large values of the wave vector or small values of the correlation length respectively. This is shown very well in Ref. [3] where we have plotted the ratio of the full characteristic frequency to its van Hove expression. This behavior is different from the nonasymptotic mode coupling expression of Olchowy [4] (a recent improvement of this theory is found in Ref. [5]) which does not yield the van Hove theory in the nonasymptotic region but instead becomes negative for large values of the variable x [3]. This region of unphysical negative values of the characteristic frequency is always reached at constant correlation length when the wave vector becomes larger than the nonuniversal parameter q_D . On the other hand the parameter q_D cannot be set to infinity as this limit yields an unphysical divergence in the hydrodynamic limit for $\xi \rightarrow 0$ [5].

4. COMPARISON WITH EXPERIMENTS

In Ref. [3] we have compared our asymptotic theory to the theoretical results of Kawasaki and Lo, Paladin and Peliti and Burstyn et al. [6] so that we shall compare our theory only to various experiments in this paper. We start this comparison with the experimental Xenon data of Swinney and Henry [7]:

In Fig. 3 we compare our asymptotic (for $c_{na} = 1$) and nonasymptotic (for $c_{na} \neq 1$) theory for the characteristic frequency ω_c/k^2 with the experimental Xenon data (all nonuniversal parameters are indicated in the figure). As discussed in Ref. [2, 3] we can treat the exponent $x_\lambda = 1 - x_\eta$ as an additional free parameter in the expression for the Onsager coefficient (see Ref. [1]) so that we can fit the initial value of the mode coupling f_0 and the exponent x_η from the experimental data for the characteristic frequency and e.g. the shear viscosity (the initial value of the Onsager coefficient Γ_0 is determined by the value of the shear viscosity at t_0). Following the fit procedure described in Ref. [1] we get the parameter f_0 from the characteristic frequency data and the exponent x_η from the microgravity shear viscosity data of Berg et al. [8]. In doing so we find good agreement for the characteristic frequency (Fig. 3) as well as for the frequency dependent shear viscosity [1].

As we can see in Fig. 3 the experimental data are not described correctly by our asymptotic expressions but only by the nonasymptotic expressions which show the crossover to the van Hove theory for large values of the reduced temperature t . Analogously any asymptotic theory [6] fails to describe the experimental data correctly. In Ref. [7] this problem was eliminated adding a regular

background contribution of the form $\omega_c^B = (\lambda^B/\rho c_p)k^2(1+x^2)$ to the critical expression for the characteristic frequency with λ^B being the regular part of the thermal conductivity and c_p the full specific heat at constant pressure containing also critical contributions. The use of the full specific heat together with the term $1+x^2$ ensures the crossover to the van Hove theory for large values of the reduced temperature as well as for large values of the wave vector (the background characteristic frequency is proportional to $k^2\xi^{-2}$ for $x \rightarrow 0$ and to k^4 for $x \rightarrow \infty$) so that the full characteristic frequency $\omega_c = \omega_c^C + \omega_c^B$ obtained by this procedure yields basically the same curves as our nonasymptotic theory (see Fig. 6 of Ref. [7]). As discussed in Ref. [3] we have to use a background characteristic frequency of the form $\omega^B = D_T^B(T, \rho)k^2 - D_T^B(T_c, \rho_c)k^2$, with the background thermal diffusivity given by $D_T^B = \lambda^B/\rho c_p^B$ and the background specific heat c_p^B containing only the regular temperature dependence without the critical singularity, in our theory. As this background term turns out to be negligibly small in the temperature range shown in Fig. 3 we have neglected it so that our asymptotic and nonasymptotic curves for Xe contain only the critical contributions discussed in this paper. So the main difference between our nonasymptotic theory and the results of Ref. [6] is that the crossover to the van Hove theory, which is clearly seen in experiments, is already contained in our expressions for the characteristic frequency and not added by an appropriate form of the background contribution!

The crossover from the asymptotic to the background behavior can be seen most easily looking at the relevant values of the mode coupling which are shown in Fig. 4. There we have plotted the mode coupling as a function of temperature

and wave vector. The experimental region (dark area) for the Xenon data covers the slope of the crossover from asymptotics to background.

CONCLUSION

We were able to show that our one-loop perturbation theory result for the characteristic frequency evaluated within the field theoretical method of the renormalization group theory does not only reproduce the correct wave vector and correlation length dependence in the hydrodynamic region as well as in the critical region but is also able to describe experimental data sufficiently well for a large range of wave vectors and reduced temperatures.

There are however some points which indicate the need for a two-loop analysis of the model: First we have seen that in one-loop order the dynamic correlation function is always of Lorentzian form whereas scaling theory [9] predicts deviations for large frequencies. Second the comparison of the frequency dependence of the shear viscosity with experimental data [1] also requires the calculation of the vertex function up to two-loop order.

ACKNOWLEDGMENTS

This work was supported by the Fonds zur Förderung der wissenschaftlichen Forschung under Project No. P12422-TPH.

References

- [1] G. Flossmann, R. Folk, and G. Moser, paper presented at the Fourteenth Symposium on Thermophysical Properties, June 25-30, 2000, Boulder, Colorado, U.S.A.
- [2] G. Flossmann, R. Folk, and G. Moser, Phys. Rev. E **60**, 779 (1999)
- [3] G. Flossmann and R. Folk, submitted to Phys. Rev. E
- [4] G. A. Olchowy and J. V. Sengers, Phys. Rev. Lett. **61**, 15 (1988)
- [5] J. Luettmer-Strathmann, J. V. Sengers, and G. A. Olchowy, J. Chem. Phys. **103**, 7482 (1995)
- [6] K. Kawasaki and S.-M. Lo, Phys. Rev. Lett. **29**, 48 (1972); G. Paladin and L. Peliti, J. Physique Lett. **43**, 15 (1982), Erratum **45**, 289 (1984); H. C. Burstyn, J. V. Sengers, J. K. Bhattacharjee, and R. A. Ferrell, Phys. Rev. A **28**, 1567 (1983)
- [7] H. L. Swinney and D. L. Henry, Phys. Rev. A **8**, 2586 (1973)
- [8] R. F. Berg, M. R. Moldover, and G. A. Zimmerli, Phys. Rev. Lett. **82**, 920 (1999); Phys. Rev. E **60**, 4079 (1999)
- [9] R. Folk and G. Moser, in *Lectures on Cooperative Phenomena in Condensed Matter Physics*, ed. D. I. Uzunov, Heron Press, 1996

FIGURE CAPTIONS

Fig. 1. Dimensionless characteristic frequency $\omega_c/\Gamma_0 k^2$ as a function of the wave vector (top) and the inverse correlation length (bottom). We can see the crossover from the asymptotics to the van Hove background.

Fig. 2. Dimensionless characteristic frequency $\omega_c/\Gamma_0 k^2$ as a function of the reduced temperature t at constant wave vector (top: $k = 10^5 \text{ cm}^{-1}$; bottom $k = 0 \text{ cm}^{-1}$) and various values of the reduced density $\Delta\rho$.

Fig. 3. Comparison of the asymptotic (dashed lines) and nonasymptotic (full lines) expressions for ω_c/k^2 with the Xe-data of Ref. [7]; from [3].

Fig. 4. Mode coupling $f_t^2(t, \omega)$ (all nonuniversal parameters for Xenon) as a function of the reduced temperature t and the dimensionless wave vector $k\xi_0$ along the critical isochore. For small values of the wave vector and the reduced temperature we reach the fixed point value $f_t^{*2} = 24/19$. The dark region marks the range of experimental data for the characteristic frequency in Xenon.




Figure 1:




Figure 2:




Figure 3:




Figure 4: